Dosing for Infants and Children





From Your Healthcare Professional

DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.

Available in:

If possible, use weight to dose; otherwise, use age. mL = milliliter

Infants' TYLENOL® Oral Suspension

Active Ingredient: Acetaminophen 160 mg (in each 5 mL)



Children's TYLENOL® Oral Suspension

Active Ingredient: Acetaminophen 160 mg (in each 5 mL)



111111111111111111111111111111111111111		SimpleMeasure*		Grape			Grape
WEIGHT	AGE	Grape Flavor USE	product only as directed.	Cherry	Cherry Florer	Use product only as directed.	Cherry
6-11 POUNDS	0-3 MONTHS	1.25 mL	2.5 mL 3.75 mL	m _L 25	_		
12-17 POUNDS	4-11 MONTHS	2.5 mL	mL 8.75 mL 5 s mL	m L 25	_		
18-23 POUNDS	12-23 MONTHS	3.75 mL	MnL	m_25	_		
24-35 POUNDS	2-3 YEARS	5 mL	2.5 m.l. m.l. 5 m.l.	1.25 m - 25	5 mL	5 mL	
36-47 POUNDS	4-5 YEARS	_			7.5 m	L 7.5 mL	
48-59 POUNDS	6-8 YEARS	_			10 ml	10 mL	
60-71 POUNDS	9-10 YEARS	_			12.5 r	mL 12.5 mL	
72-95 POUNDS	11 YEARS	_			15 ml	- 15 mL	

Kids' Wellness Tracker

This free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN® makes it easy to track your child's height, weight, BMI, vaccines, symptoms, and medicines, plus calculate dosing.

Available on the App Store[™] and the Google Play Store[™]





Google Play is a trademark of Google Inc. App Store is a service mark of Apple Inc. Standard download rates may apply.

IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: _____

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all TYLENOL® products.
- Repeat dose every 4 hours while symptoms last.
- Do NOT give more than 5 doses in 24 hours.
- Do NOT use with any other product containing acetaminophen.
- Use only the dosing device (syringe or dosing cup) that came with the product.
- All Infants' TYLENOL® and Children's TYLENOL® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL).

Dosing for Infants and Children

From Your Healthcare Professional





DOSE: Every 6-8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

Available in:

Motrin If possible, use weight to dose; otherwise, use age. Lasts up to 8 hours **mL** = milliliter

Infants' MOTRIN® **Concentrated Drops**

Active Ingredient:

Ibuprofen 50 mg (NSAID)* (in each 1.25 mL)

*Nonsteroidal anti-inflammatory drug



Children's MOTRIN® **Oral Suspension**

Active Ingredient: Ibuprofen 100 mg (NSAID)*

(in each 5 mL)

*Nonsteroidal anti-inflammatory drug





		3 0	Dye-Free	
WEIGHT	AGE	Be	Dye-Free Berry Berry Use product only as directed. Dye-Berry Berry Dye-Berry Dye-Berry	У
_	<6 MONTHS	_	_	
12-17 POUNDS	6-11 MONTHS	1.25 mL		
18-23 POUNDS	12-23 MONTHS	1.875 mL		
24-35 POUNDS	2-3 YEARS	_	5 mL 5 mL	
36-47 POUNDS	4-5 YEARS	_	7.5 mL	
48-59 POUNDS	6-8 YEARS	_	10 mL	
60-71 POUNDS	9-10 YEARS	_	12.5 mL	
72-95 POUNDS	11 YEARS	_	15 mL	

Kids' Wellness Tracker

This free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN® makes it easy to track your child's height, weight, BMI, vaccines, symptoms, and medicines, plus calculate dosing.

Available on the App Store[™] and the Google Play Store $^{\text{\tiny{M}}}$





Google Play is a trademark of Google Inc. App Store is a service mark of Apple Inc. Standard download rates may apply.

IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: _

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all MOTRIN® products.
- Repeat dose every 6-8 hours as needed.
- Do NOT give more than 4 doses in 24 hours.
- Use only the dosing device (syringe or dosing cup) that came with the product.